

Electromagnetic Form Factors of Hadrons in Dual-Large N_c QCD ¹

1

C. A. Dominguez

Centre for Theoretical Physics and Astrophysics
University of Cape Town, Rondebosch 7700, South Africa, and Department of Physics,
Stellenbosch University, Stellenbosch 7600, South Africa

Abstract

In this talk, results are presented of determinations of electromagnetic form factors of hadrons (pion, proton, and $\Delta(1236)$) in the framework of Dual-Large N_c QCD (Dual- QCD_∞). This framework improves considerably tree-level VMD results by incorporating an infinite number of zero-width resonances, with masses and couplings fixed by the dual-resonance (Veneziano-type) model.

¹Invited talk at the XII Mexican Workshop on Particles & Fields, Mazatlan, November 2009. To be published in American Institute of Physics Conference Proceedings Series. Work supported in part by NRF (South Africa).

1 Introduction

In its original formulation [1], Vector Meson Dominance (VMD) is an effective tree-level model based on $\gamma - \rho^0$ conversion. If used to compute electromagnetic form factors, it can very roughly account for the pion form factor data in the space-like region, and with some modifications (unitarization), also in the time-like region around the rho-meson peak. However, for non-zero spin hadrons such as nucleons and $\Delta(1236)$, VMD is in serious disagreement with the observed q^2 fall-off of these form factors. It has been generally believed that the reason for this discrepancy is that the radial excitations of the ρ -meson are not taken into account in naive VMD. In fact, an infinite sum of monopole terms with suitable coefficients can lead to a form factor with an asymptotic behavior different from a monopole. An attempt to remedy this situation was made long ago by incorporating radial excitations of the rho-meson into VMD, i.e. Extended VMD [2]. At the time, though, there was no known renormalizable QFT to support this approach. Today, we know that in the limit of an infinite number of colors, QCD is solvable and leads to a hadronic spectrum consisting of an infinite number of zero-width states [3]. However, the masses and couplings of these states remain unspecified, so that models are needed to fix these parameters. An attractive and highly economical candidate is Dual- QCD_∞ [4]-[7], **inspired** in the Dual Resonance Model for scattering amplitudes of Veneziano [8], the precursor of string theory. It is very important to stress the word **inspired**, as Dual- QCD_∞ does not share any of the unwanted features of the original Veneziano model for n-point functions ($n \geq 4$), such as lack of unitarity, unphysical particles (tachyons) in the spectrum, etc. These features simply do not emerge for three-point functions. Another aspect of Dual- QCD_∞ which needs to be stressed to avoid misunderstandings is that it is not intended to be an expansion in powers of $1/N_c$. In fact, N_c is taken to be infinite from the start, as this is the limit in which QCD is solvable and leads to the hadronic spectrum mentioned above. Unitarization can subsequently be performed by shifting the poles from the real axis into the second Riemann sheet in the complex energy (squared) plane. This induces corrections to form factors of order $\mathcal{O}(\Gamma/M \simeq 10\%)$.

2 Dual- QCD_∞

In QCD_∞ , a typical form factor has the generic form

$$F(s) = \sum_{n=0}^{\infty} \frac{C_n}{(M_n^2 - s)}, \quad (1)$$

where $s \equiv q^2$ is the momentum transfer squared, and the masses M_n , and the couplings C_n remain unspecified. In Dual- QCD_∞ they are given by [4]

$$C_n = \frac{\Gamma(\beta - 1/2)}{\alpha' \sqrt{\pi}} \frac{(-1)^n}{\Gamma(n+1)} \frac{1}{\Gamma(\beta - 1 - n)}, \quad (2)$$

where β is a free parameter, and the string tension α' is $\alpha' = 1/2M_\rho^2$, as it enters the rho-meson Regge trajectory $\alpha_\rho(s) = 1 + \alpha'(s - M_\rho^2)$. The mass spectrum is chosen as $M_n^2 = M_\rho^2(1 + 2n)$. This simple formula correctly predicts the first few radial excitations. Other, e.g. non-linear




Figure 1: Pion form factor in Dual- QCD_∞ (solid curve), and VMD (broken curve).




Figure 2: Pion form factor in Dual- QCD_∞ in the time-like region.




Figure 3: Sachs magnetic form factor of the proton, $G_M(Q^2)$. Data is from the reanalysis of [10], and $Q^2 \equiv -q^2$.

mass formulas could be used, but this hardly changes the results in the space-like region, and only affects the time-like region behavior for very large q^2 . With these choices the form factor becomes an Euler Beta-function, i.e.

$$\begin{aligned}
 F(s) &= \frac{\Gamma(\beta - 1/2)}{\sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n}{\Gamma(n+1)} \frac{1}{\Gamma(\beta - 1 - n)} \frac{1}{[n + 1 - \alpha_\rho(s)]} \\
 &= \frac{1}{\sqrt{\pi}} \frac{\Gamma(\beta - 1/2)}{\Gamma(\beta - 1)} B(\beta - 1, 1/2 - \alpha' s),
 \end{aligned} \tag{3}$$

where $B(x, y) = \Gamma(x)\Gamma(y)/\Gamma(x+y)$. The form factor exhibits asymptotic power behavior in the space-like region, i.e.

$$\lim_{s \rightarrow -\infty} F(s) = (-\alpha' s)^{(1-\beta)}, \tag{4}$$

from which one identifies the free parameter β as controlling this asymptotic behavior. Notice that while each term in Eq.(3) is of the monopole form, the result is not necessarily of this form because it involves a sum over an infinite number of states. The exception occurs for integer values of β , which leads to a finite sum. The imaginary part of the form factor Eq.(3) is

$$\text{Im } F(s) = \frac{\Gamma(\beta - 1/2)}{\alpha' \sqrt{\pi}} \sum_{n=0}^{\infty} \frac{(-1)^n}{\Gamma(n+1)} \frac{1}{\Gamma(\beta - 1 - n)} \pi \delta(M_n^2 - s). \tag{5}$$

Unitarization can be performed by shifting the poles from the real axis in the complex s -plane. The simplest model is the Breit-Wigner form

$$\pi \delta(M_n^2 - s) \rightarrow \frac{\Gamma_n M_n}{[(M_n^2 - s)^2 + \Gamma_n^2 M_n^2]}, \tag{6}$$

where one expects Γ_n to grow with M_n . Other, more refined choices, are certainly possible, e.g. the Gounaris-Sakurai form in which the width is momentum transfer dependent.




Figure 4: Ratio of electric to magnetic Sachs form factors of the proton. Data is from the reanalysis of [10], and $Q^2 \equiv -q^2$.




Figure 5: $\Delta(1236)$ magnetic form factor in Dual- QCD_∞ (solid line).

Results in this framework are shown, together with the data, in Fig. 1 and Fig.2 for the pion (data from [9]), Figs. 3 and 4 for the proton, and Fig. 5 for the $\Delta(1236)$. The time-like pion form factor was obtained using Eq.(6); in spite of the simplicity of the model the agreement is reasonable at and around the ρ -peak. In the case of the proton, Eq.(3) is used for the Dirac and Pauli form factors, $F_1(q^2)$ and $F_2(q^2)$, as these have the correct analyticity properties. The fits have not been made to the raw data, but rather to the data base as corrected in [10]. These corrections take into account the discrepancies between unpolarized (SLAC) and polarized (JLAB) experiments. For the $\Delta(1236)$, the three so called Scadron form factors $G_{M,E,C}^*(q^2)$ were fitted using Eq.(3), and data on $G_M^*(q^2)$, and the two ratios between $G_{E,C}^*(q^2)$ and $G_M^*(q^2)$ [11]-[12]. The value of the free parameter β in the form factor, Eq. (3), which determines its asymptotic behavior, is as follows: for the pion, $\beta_\pi = 2.3$, for F_1 and F_2 of the proton, $\beta_1 = 2.95 - 3.03$, and $\beta_2 = 4.13 - 4.20$, respectively, and for G_M^* , G_E^* , G_C^* of the $\Delta(1236)$, $\beta_M^* = 4.6 - 4.8$, $\beta_E^* \simeq \beta_M^*$, and $\beta_C^* = 6.0 - 6.2$. Taking the middle values of these numbers, the asymptotic behavior in the space-like region of these form factors is approximately as follows: $F_\pi \sim (-q^2)^{-1.3}$, $F_1 \sim (-q^2)^{-2.0}$, $F_2 \sim (-q^2)^{-3.2}$, $G_M^* \sim G_E^* \sim (-q^2)^{-3.7}$, and $G_C^* \sim (-q^2)^{-5.1}$.

3 Conclusions

Dual- QCD_∞ accounts quite successfully for the space-like behavior of the pion, the proton, and the $\Delta(1236)$ electromagnetic form factors. A very simple unitarization procedure leads to a pion form factor in the time-like region in reasonable agreement with data (more sophisticated procedures improve considerably this agreement [6]). Nucleon form factors in this framework and in the time-like region are currently under investigation [13]

4 Acknowledgments

This talk is based on work done by the author, and in collaboration with R. Röntsch, and T. Thapedi. Work supported in part by the NRF (South Africa). The author thanks the organizers for an interesting and fruitful workshop.

References

- [1] J. J. Sakurai, *Currents and Mesons*, University of Chicago Press, Chicago, 1969.
- [2] A. Bramon, E. E. Etim, and M. Greco, *Phys. Lett.* **B 41**, 607 (1972); M. Greco, *Nucl. Phys.* **B 63**, 398 (1973).
- [3] G. 't Hooft, *Nucl. Phys.* **B 72**, 461 (1974); E. Witten, *Nucl. Phys.* **B 79**, 57 (1979).
- [4] C. A. Dominguez, *Phys. Lett.* **B 512**, 331 (2001).

- [5] C. A. Dominguez, and T. Thapedi, *J. High Energy Phys.* **0410**, 003 (2004).
- [6] C. Bruch, A. Khodjamirian, J. H. Kühn, *Eur. Phys. J. C* **39**, 41 (2005).
- [7] C. A. Dominguez, and R. Röntsch, *J. High Energy Phys.* **0710**, 085 (2007).
- [8] P. H. Frampton, *Dual Resonance Models*, Benjamin, Reading, Massachusetts, 1974.
- [9] C. J. Bebek et al., *Phys. Rev. D* **17**, 1693 (1978); P. Brauel et al., *Z. Phys. C* **3**, 101 (1979); J. Volmer et al., *arXiv: Nucl-ex /0010009* (2000); S. R. Amendolia et al., *Nucl.Phys. B* **277**, 168 (1986).
- [10] E. J. Brash, A. Kozlov, Sh. Li, G. M. Huber, *Phys. Rev. C* **65**, 051001 (2002).
- [11] W. Bartel et al., *Phys. Lett. B* **28**, 148 (1968); J. Bleckwien et al., *DESY Report 71/63* (1971), unpublished; K. Bätzner et al., *Phys. Lett. B* **39**, 575 (1972); J. C. Alder et al., *Nucl. Phys. B* **46**, 573 (1972); S. Stein et al., *Phys. Rev. D* **12**, 1884 (1975); P. Stoler, Phys. Rept. **226**, 103 (1993); L. M. Stuart et al., *Phys. Rev. D* **58**, 032003 (1998).
- [12] V. V. Frolov et al., *Phys. Rev. Lett.* **82**, 45 (1999) ; R. Beck et al., *Phys. Rev. C* **61**, 035204 (2000); K. Joo et al., Phys. Rev. Lett. **88**, 122001 (2002); N. F. Sparveris, *Phys. Rev. Lett.* **94**, 022003 (2005); J. J. Kelly et al., Phys. Rev. Lett. **95**, 102001 (2005); M. Ungaro et al., Phys. Rev. Lett. **97**, 112003 (2006); S. Stave et al., Eur. Phys. J. **A 30**, 471 (2006).
- [13] C.A. Dominguez (work in progress).